arXiv:1304.5832v1 [math-ph] 22 Apr 2013

On the marginally trapped surfaces in Minkowski space-time with finite type Gauss map

Nurettin Cenk Turgay*

April 23, 2013

Abstract

In this paper, we work on the marginally trapped surfaces in the Minkowski space-time $\mathbb{E}^4_1$. We obtain the complete classification of the marginally trapped surfaces with pointwise 1-type Gauss map. We also proved that if a marginally trapped surface has non-harmonic, 1-type Gauss map, then it is null 2-type. Further, we give construction of marginally trapped surfaces with 1-type Gauss map with a given boundary curve. We also state some explicit examples.

Keywords: Minkowski space-time, marginally trapped surface, finite type Gauss map, null 2-type Mathematics Subject Classification.: 53B25, 53C40

1 Introductions

Let M be an n-dimensional semi-Riemannian submanifold of a semi-Euclidean space and ϕ be a mapping defined on M into another semi-Euclidean space. Considering the spectral decomposition of ϕ with respect to the Laplace operator Δ of M is one of the natural ways to study ϕ. Moreover, if ϕ is one of geometrically important mappings on M such as its Gauss map or its position vector, then understanding spectrum of ϕ may play great role to understand geometry of M. In this direction, the notion of finite type mappings was introduced by B. Y. Chen in late 1970's. Many works have been published in this topic, [3, 4, 5, 6, 8, 9].

By the definition, if the mapping ϕ can be expressed as a sum of finitely many eigenvectors of Δ, then it is said to be of finite type. More precisely,

---

*e-mail:turgayn@itu.edu.tr, Adress:Istanbul Technical University, Faculty of Science and Letters, Department of Mathematics, 34469 Maslak, Istanbul, Turkey

if these eigenvectors are corresponding from k distinct eigenvalues of Δ, then ϕ is said to be of k-type. Some important results about finite type mappings defined on semi-Riemannian submanifolds have appeared sofar (cf. [1, 8, 20]).

Particularly, the Gauss map of submanifolds has been worked in several articles in terms of being finite type after some results on the submanifold with 1-type Gauss map or 2-type Gauss map had been given in [9]. The Gauss map ν of the submanifold M is 1-type if and only if it satisfies

$$\Delta\nu = \lambda(\nu + C) \tag{1.1}$$

for some $\lambda \in \mathbb{R}$ and some constant vector C. However, there are some geometrically important submanifolds whose Gauss map satisfies a weaker condition. It is well-known that the Laplacian of the Gauss map of surfaces such as helicoids and catenoids in the 3-dimensional Minkowski space-time and generalized torus in the 4-dimensional Euclid space takes the form of

$$\Delta\nu = f(\nu + C) \tag{1.2}$$

for some smooth function f on M and some constant vector C. A submanifold with the Gauss map satisfying (1.2) is said to have pointwise 1-type Gauss map (cf. [12, 13, 17, 22]).

On the other hand, marginally trapped surfaces play an important role in the concept of trapped surfaces, introduced by Penrose in [18]. Let M be a space-like surface in the Minkowski space-time and H its mean curvature vector. In the aspect of general relativity, H measures the tension of the surface coming from the surrounding space. M is said to be trapped if H is time-like on M. In this case, if H is future- or past-pointing, then M is called future- or past-trapped surface, respectively. A trapped surface has physical property that the outgoing light rays from it are convergent. It is believed that there is a surface, called marginally trapped (or quasi-minimal) surface, to which all the light rays are parallel. Furthermore, a marginally trapped surface locates the black hole, [2, 10, 16].

Geometrically, a space-like surface in a 4-dimensional space-time is called marginally trapped if its mean curvature vector is light-like. In the very recent past, many problems in geometry involving marginally trapped surfaces in a space-time have been dealed with (cf. [2, 7, 10, 11, 16]). For example, complete classification of biharmonic marginally trapped surfaces was obtained in [7]. Haesen and Ortega classified marginally trapped surfaces which are invariant under a group of boost isometries in [16]. Recently, in [2], isotropic marginally trapped surfaces were classified by Cabrerizo et al. In

[21], a generalization of trapped surfaces was obtained for the submanifolds of a Lorentzian space with codimension 2.

In this article, we study on the Gauss map of marginally trapped surfaces in the Minkowski space-time. In Section 2, after we describe our notations, which are along the lines of notations used in [10], we give a summary of the basic facts and formulas that we will use. In Section 3, we give a characterization of marginally trapped surfaces with pointwise 1-type Gauss map of the first kind. In Section 4, we construct a class of pseudo-umbilical marginally trapped surfaces with pointwise 1-type Gauss map of the second kind. In Section 5, first, we prove that if the Gauss map of a marginally trapped surface in the Minkowski space-time satisfies (1.2) for $C \neq 0$ and $f \not\equiv 0$, then it must be a member of the family of pseudo-umbilical marginally trapped surfaces that we have construced in Section 4. Then, we complete classification of marginally trapped surfaces with pointwise 1-type Gauss map. Further, we give some methods for construction of marginally trapped surfaces with 1-type Gauss map. We also give some explicit examples.

The surfaces we are dealing with are smooth and connected unless otherwise stated.

2 Preliminaries

2.1 Basic notation, formulas and definitions

Let $\mathbb{E}^m_s$ denote the pseudo-Euclidean m-space with the canonical pseudo-Euclidean metric tensor g of index s given by

$$g = -\sum_{i=1}^{s} dx_i^2 + \sum_{j=s+1}^{m} dx_j^2,$$

where $(x_1, x_2, \ldots, x_m)$ is a rectangular coordinate system in $\mathbb{E}^m_s$. We put

$$\begin{aligned} \mathbb{S}^{m-1}_s(r^2) &= \{x \in \mathbb{E}^m_s : \langle x, x \rangle = r^{-2}\}, \\ \mathbb{H}^{m-1}_{s-1}(-r^2) &= \{x \in \mathbb{E}^m_s : \langle x, x \rangle = -r^{-2}\}, \end{aligned}$$

where $\langle\ ,\ \rangle$ is the indefinite inner product of $\mathbb{E}^m_s$. In general relativity, $\mathbb{E}^4_1$, $\mathbb{S}^4_1(r^2)$ and $\mathbb{H}^4_1(r^2)$ are known as the Minkowski, de Sitter and anti-de Sitter space-times, respectively, [11].

A vector $\zeta \neq 0 \in T_p(\mathbb{E}^m_s) \cong \mathbb{E}^m_s$ is called space-like (resp. time-like or light-like) if $\langle \zeta, \zeta \rangle > 0$ (resp. $\langle \zeta, \zeta \rangle < 0$ or $\langle \zeta, \zeta \rangle = 0$), where $T_p(\mathbb{E}^m_s)$ denotes the tangent space of $\mathbb{E}^m_s$ at p.

Let M be an n-dimensional immersed semi-Riemannian submanifold of the semi-Euclidean space $\mathbb{E}^m_s$ and Levi-Civita connections of $\mathbb{E}^m_s$ and M be denoted by $\widetilde{\nabla}$ and ∇, respectively.

In this section, we shall use letters X, Y, (resp. ξ, η) to denote vectors fields tangent (resp. normal) to M. The Gauss and Weingarten formulas are given by

$$\begin{aligned}\widetilde{\nabla}_X Y &= \nabla_X Y + h(X,Y) \qquad (2.1)\\ \widetilde{\nabla}_X \xi &= -A_\xi X + D_X \xi, \qquad (2.2)\end{aligned}$$

where h, D and A are the second fundamental form, the normal connection and the shape operator of M, respectively. On the other hand, the shape operator A and the second fundamental form h of M are related by

$$\langle A_\xi X, Y\rangle = \langle h(X,Y), \xi\rangle. \qquad (2.3)$$

The mean curvature vector of M is defined by $H = \frac{1}{n}\mathrm{tr}h$.

Denote by R and R^D the curvature tensor associated with the connections ∇ and D, respectively. Then, the Ricci equation is given by

$$\langle R^D(X,Y)\xi, \eta\rangle = \langle [A_\xi, A_\eta]X, Y\rangle. \qquad (2.4)$$

Let M be a surface in the Minkowski space-time $\mathbb{E}^4_1$. M is said to be space-like if every non-zero tangent vector on M is space-like. Consider an orthonormal frame field $\{e_1, e_2, e_3, e_4\}$ on M. The connection forms ω_{AB} associated with this frame field are defined by $\omega_{AB}(X) = \langle \widetilde{\nabla}_X e_A, e_B\rangle$ and satisfy $\omega_{AB} + \omega_{BA} = 0$ for $A, B = 1, 2, 3, 4$.

The Gaussian curvature K of M is defined by $K = R(e_1, e_2, e_2, e_1)$. A surface is said to be flat if $K \equiv 0$. The normal curvature K^D of M is defined by $K^D = R^D(e_1, e_2; e_3, e_4)$. If K^D vanishes identically, M is said to have flat normal bundle.

Let the mean curvature vector H of M satisfies $\langle H, H\rangle \equiv 0$ on M. If H is light-like on M then, it is said to be a marginally trapped surface. If H vanishes on a part of M, then the surface is called partly marginally trapped, [16].

Throughout this work we denote the norm of e_A by ε_A, i.e., $\varepsilon_A = \langle e_A, e_A\rangle = \mp 1$. In addition, the subscripts u, v etc. denotes the partial derivatives with respect to u, v etc.

2.2 Gauss map

Let M be an n-dimensional, oriented semi-Riemannian submanifold of the semi-Euclidean space $\mathbb{E}^m_s$. We choose a local, orthonormal frame field $\{e_1, e_2, \ldots, e_m\}$

defined on M, where $e_1,\ e_2,\ \ldots,\ e_n$ (resp. $e_{n+1},\ e_{n+2},\ \ldots,\ e_m$) are tangent (resp. normal) to M. Then, the Laplace operator of M with respect to the induced metric is defined by

$$\Delta = \sum_{i=1}^{n} \varepsilon_i(-e_i e_i + \nabla_{e_i} e_i).$$

Let $\mathbb{E}_s^m$ and $G(m-n,m)$ denote the space of $(m-n)$-vectors on $\mathbb{E}_s^m$ and the Grassmannian manifold consisting of all $(m-n)$-planes through the origin of $\mathbb{E}_s^m$, respectively. Note that $G(m-n,m)$ is canonically imbedded in $\Lambda^{m-n}(\mathbb{E}_s^m)$ and $\Lambda^{m-n}(\mathbb{E}_s^m)$ is an N dimensional vector field, where $N = \binom{m}{m-n}$. A non-degenerate inner product on $\Lambda^{m-n}(\mathbb{E}_s^m)$ is defined by

$$\langle X_1 \wedge X_2 \wedge \cdots \wedge X_{m-n}, Y_1 \wedge Y_2 \wedge \cdots \wedge Y_{m-n} \rangle = \det(\langle X_i, Y_j \rangle),$$

where $X_i,\ Y_i \in \mathbb{E}_s^m,\ i = 1,2,\ldots,m-n$ and $\langle X_i, Y_j \rangle$ denotes the semi-Euclidean inner product of the vectors X_i and Y_j. We will denote the inner product space $\left(\Lambda^{m-n}(\mathbb{E}_s^m), \langle , \rangle\right)$ by $\Lambda^{m,n}$.

Note that, the set $\mathcal{B} = \{(e_{i_1} \wedge e_{i_2} \wedge \ldots \wedge e_{i_{m-n}})(p) |\ 1 \leq i_1 < i_2 < \cdots < i_n \leq m\}$ is an orthonormal basis for the space $\Lambda^{m,n}$. Let S be the number of time-like vectors in $\mathcal{B}$. Then, there exist a one-to-one, onto and linear isometry from $\Lambda^{m,n}$ into $\mathbb{E}_S^N$, because their dimension and index are equal(see [19, p. 52]). Hence, we have $G(m-n,m) \subset \Lambda^{m,n} \cong \mathbb{E}_s^m$. Therefore, one can assume the $(m-n)$-vector $X_1 \wedge X_2 \wedge \ldots \wedge X_{m-n}$ as a vector in $\mathbb{E}_S^N$.

The smooth mapping

$$\begin{array}{rcl} \nu : M & \rightarrow & G(m-n,m) \subset M_S^{N-1}(\varepsilon) \subset \mathbb{E}_S^N \cong \Lambda^{m,n} \\ p & \mapsto & \nu(p) = (e_{n+1} \wedge e_{n+2} \wedge \ldots \wedge e_m)(p) \end{array} \tag{2.5}$$

is called the (normal) Gauss map of M which assigns to a point p in M the oriented $(m-n)$-plane through the origin of $\mathbb{E}_s^m$ and parallel to the normal space of M at p, where

$$\widetilde{M}_S^{N-1}(\varepsilon) = \begin{cases} \mathbb{S}_S^{N-1}(1) & \text{in } \mathbb{E}_S^N \quad \text{if } \varepsilon = 1 \\ \mathbb{H}_{S-1}^{N-1}(-1) & \text{in } \mathbb{E}_S^N \quad \text{if } \varepsilon = -1 \end{cases}$$

and $\varepsilon = \langle \nu, \nu \rangle = \varepsilon_{n+1}\varepsilon_{n+2} \cdots \varepsilon_m = \pm 1$, [17].

A semi-Riemannian submanifold M of a semi-Euclidean space $\mathbb{E}_s^m$ is said to have harmonic Gauss map if $\Delta\nu \equiv 0$ on M. On the other hand, M is said to have *pointwise 1-type Gauss map* if it satisfies (1.2) for a constant vector $C \in \mathbb{E}_S^N$ and a smooth fuction $f \not\equiv 0$. Moreover, a pointwise 1-type Gauss map is called *of the first kind* if (1.2) is satisfied $C = 0$, and *of the second kind* if $C \neq 0$. Moreover, if (1.2) is satisfied for a non-constant function f, then M is said to have *proper* pointwise 1-type Gauss map.

2.3 Eigenvalues of Laplace equation in the plane

In this subsection, we use the same notation with [15] and review briefly some known facts on the eigenvalues of Laplace operator given by

$$\begin{array}{rcl} \Delta : H_0^1(\Omega) & \rightarrow & H_0^1(\Omega) \\ \phi & \mapsto & -\phi_{uu} - \phi_{vv}, \end{array} \tag{2.6}$$

where Ω is an open bounded subset of $\mathbb{R}^2$ with the boundary curve $\beta = \partial\Omega$.

Note that a function ϕ is an eigenfunction of Δ if and only if it is a solution of boundary value problem

$$\left\{ \begin{array}{rcl} \Delta\phi(u,v) + \lambda\phi(u,v) & = & 0 \quad \text{if } (u,v) \in \Omega \\ \phi & = & 0 \quad \text{if } (u,v) \in \beta. \end{array} \right. \tag{2.7}$$

As Δ is an elliptic, symmetric operator, the following arguments hold (see [15, p. 326-335]). The eigenvalues of Δ are $0 < \lambda_1 \leq \lambda_2 \leq \ldots \nearrow \infty$. Moreover, there exists an orthonormal basis $\{\phi_k | k = 1, 2 \ldots\}$ of $L^2(\Omega)$ where $\phi_k \in H_0^1(\Omega)$ and satisfies (2.7) for $\lambda = \lambda_1, \lambda_2, \ldots$. The regularity theory shows that $\phi \in C^\infty(\Omega)$. Moreover, if β is smooth, then $\phi \in C^\infty(\bar{\Omega})$.

3 Marginally trapped surfaces with pointwise 1-type Gauss map of the first kind

In [14], the author and U. Dursun obtained characterization of space-like surfaces in the Minkowski space-time with pointwise 1-type Gauss map of the first kind. In this section, we give a short brief of the results on marginally trapped surfaces obtained in [14].

First, we obtain the Laplacian of the Gauss map in the following lemma.

Lemma 3.1. *Let M be an oriented, partly marginally trapped surface in the Minkowski space-time and $\{e_1, e_2\}$ be a local orthonormal base field of the tangent bundle of M. Then, the Laplacian of the Gauss map ν given by* (2.5) *is*

$$\Delta\nu = -2K\nu + 2K^D e_1 \wedge e_2 + 2\sum_{i=1}^{2} D_{e_i}H \wedge e_i, \tag{3.1}$$

where K, K^D and H are the Gaussian curvature, the normal curvature and the mean curvature vector of M, respectively.

It is well-known that for a space-like surface M in $\mathbb{E}^4_1$, being parallel of H implies $K^D \equiv 0$. Thus, the next corollary directly follow from Lemma 3.1.

Corollary 3.2. *[14] Let M be a marginally trapped surface in the Minkowski space-time. Then, M has pointwise 1-type Gauss map of the first kind if and only if M has parallel mean curvature vector. Moreover, M has harmonic Gauss map if and only if M is a flat surface with parallel mean curvature vector.*

Remark 1. See [11], for the classification of flat marginally trapped surfaces in the Minkowski space-time with parallel mean curvature vector.

In [11], the classification of marginally trapped surfaces in the Minkowski space-time with parallel mean curvature vector is given. By combaining [11, Theorem 4.1] and Corollary 3.2, we state

Proposition 3.3. *[14] Let M be a marginally trapped surface in the Minkowski space-time $\mathbb{E}^4_1$. Then, M has proper pointwise 1-type Gauss map of the first kind if and only if it is congruent to one of the following 2 type of surfaces:*

(i) a non-flat, non-parallel surface lying in the de Sitter space-time $S^3_1(r^2)$ for some $r > 0$ such that the mean curvature vector H' of M in $S^3_1(r^2)$ satisfies $\langle H', H'\rangle = -r^2$;

(ii) a non-flat, non-parallel surface lying in the hyperbolic space $H^3(-r^2)$ for some $r > 0$ such that the mean curvature vector H' of M in $H^3(-r^2)$ satisfies $\langle H', H'\rangle = r^2$.

Proof. Let M be a a marginally trapped surface in the Minkowski space-time with proper pointwise 1-type Gauss map of the first kind. Then, M is non-flat and it has parallel mean curvature vector, because of Corollary 3.2. According to [11, Theorem 4.1], a non-flat marginally trapped surface with parallel mean curvature vector is congruent to one of the surfaces given in the proposition.

Conversely, if we assume M is congruent to one of the surfaces given in the proposition, then Lemma 3.1 implies $\Delta\nu = -2K\nu$, where K is the Gaussian curvature of M. Thus, M has pointwise 1-type Gauss map of the first kind. Moreover, by a computation using Codazzi and Gauss equations, one can see that if the Gaussian curvature of a surface lying in $S^3_1(r^2)$ or $H^3(-r^2)$ is constant, then it must be zero (see [14] for the details). Thus, $K \not\equiv$ implies that K is not constant. $\square$

4 Gauss map of pseudo-umbilical marginally trapped surfaces in $\mathbb{E}^4_1$

A submanifold in $\mathbb{E}^m_s$ is said to be pseudo-umbilical if there exists a function ρ such that the second fundamental form h and the mean curvature vector H of M satisfy

$$\langle h(X,Y),H\rangle=\rho\langle X,Y\rangle \tag{4.1}$$

for all tangent vector fields X,Y of M. In this case, (4.1) is satisfied for the function ρ given by $\rho=\langle H,H\rangle$, [2]. Thus, from (2.3) and (4.1) one can see that a marginally trapped surface in $\mathbb{E}^4_1$ is pseudo-umbilical if and only if the shape operator A_H along the mean curvature vector of M is a zero-operator.

Remark 2. Let M be a (partly) marginally trapped surface in $\mathbb{E}^4_1$. If $A_H\equiv 0$ on M, then M is congruent to the surface given by

$$x(u,v)=(\phi(u,v),u,v,\phi(u,v)), \tag{4.2}$$

for a smooth function $\phi:\Omega\to\mathbb{R}$, where Ω is an open subset of $\mathbb{R}^2$ (see the proof of Theorem 6.1 in [7] and also [2]).

Let M be the surface in $\mathbb{E}^4_1$ given by (4.2) for a smooth function $\phi:\Omega\to\mathbb{R}$. We choose following moving orthonormal frame field $\{e_1,e_2,e_3,e_4\}$ such that

$$e_1=\frac{\partial}{\partial u},\quad e_2=\frac{\partial}{\partial v}, \tag{4.3a}$$

$$e_3=\frac{1}{\xi}(0,-\phi_u,-\phi_v,1), \tag{4.3b}$$

$$e_4=-\frac{1}{\xi}(\phi_u^2+\phi_v^2+1,\phi_u,\phi_v,\phi_u^2+\phi_v^2), \tag{4.3c}$$

where $\xi=\sqrt{\phi_u^2+\phi_v^2+1}$. Note that from (4.3b) and (4.3c) we have

$$e_3-e_4=\xi(1,0,0,1). \tag{4.4}$$

By a direct computation, we obtain $\omega_{12}=0$ and

$$A_3=A_4=\frac{1}{\xi}\begin{pmatrix}\phi_{uu} & \phi_{uv}\\ \phi_{uv} & \phi_{vv}\end{pmatrix}. \tag{4.5}$$

Thus, $K\equiv 0$ and (2.4) implies $K^D\equiv 0$ on M. Moreover, the mean curvature vector H becomes $H=-\frac{\Delta\phi}{2}(1,0,0,1)$ because of (4.4). Thus, from (3.1) we obtain

$$\Delta\nu=-(1,0,0,1)\wedge\left(\Delta\phi_u e_1+\Delta\phi_v e_2\right). \tag{4.6}$$

On the other hand, from (4.3b) and (4.4) we have

$$\nu = (1,0,0,1) \wedge (\xi e_3) = (1,0,0,1) \wedge (0, -\phi_u, -\phi_v, 1). \tag{4.7}$$

In the next proposition, we obtain a family of pseudo-umbilical surfaces with pointwise 1-type Gauss map of the second kind.

Proposition 4.1. *Let* Ω *be an open, bounded subset of* $\mathbb{R}^2$, $\psi, \phi : \Omega \to \mathbb{R}$ *some smooth functions satisfying*

$$\Delta\psi = F(\psi), \tag{4.8a}$$

$$\phi(u,v) = \psi(u,v) + c_1 u + c_2 v \tag{4.8b}$$

for some constants c_1, c_2 *and a differentiable non-constant function* $F : \psi(\Omega) \to \mathbb{R}$ *such that the function* $f : \Omega \to \mathbb{R}$ *defined by*

$$f(u,v) \quad = \quad F'(\psi(u,v)) \tag{4.9}$$

is smooth. Assume that M *is the partly marginally trapped surface in the Minkowski space-time given by* (4.2).

Then, the Gauss map of M *is pointwise 1-type of the second kind and satisfies* (1.2) *for the smooth function* f *given in* (4.9) *and the constant vector*

$$C \quad = \quad -(1,0,0,1) \wedge (0, c_1, c_2, 1). \tag{4.10}$$

Proof. Let $\{e_1, e_2, e_3, e_4\}$ be the frame field on M given by (4.3). If ϕ satisfies (4.8), then the equations (4.6), (4.8) and (4.9) imply $\Delta\nu = -f(1,0,0,1) \wedge \Big((\phi_u + c_1)e_1 + (\phi_v + c_2)e_2\Big)$ from which and (4.3a) we obtain

$$\begin{aligned} \Delta\nu =& f(1,0,0,1) \wedge (0, \phi_u + c_1, \phi_v + c_2, 0) \\ =& f\Big((1,0,0,1) \wedge (0, -\phi_u, -\phi_v, 1) - (1,0,0,1) \wedge (0, c_1, c_2, 1)\Big). \end{aligned}$$

From this equation, (4.7) and (4.10) we obtain (1.2). Hence, M has pointwise 1-type Gauss map of the second kind. ☐

Next, we obtain the following classification theorem of pseudo-umbilical marginally trapped surfaces in $\mathbb{E}^4_1$ with pointwise 1-type Gauss map.

Proposition 4.2. *Let* M *be a pseudo-umbilical marginally trapped surface in the Minkowski space-time. Then,* M *has pointwise 1-type Gauss map of the second kind if and only if* M *is congruent to a surface given in Proposition 4.1.*

Proof. Let M be a marginally trapped surface in the Minkowski space-time given by (4.2). We consider the frame field $\{e_1, e_2, e_3, e_4\}$ on M given by (4.3). Then, the Gauss map ν of M satisfies (4.6) and (4.7).

Now, in order to prove the necessary condition, we assume that M has pointwise 1-type Gauss map of the second kind. Then, the equation (1.2) is satisfied for a smooth function $f \not\equiv 0$ and a constant vector $C \neq 0$.

If $f \equiv 0$ on an open, connected subset $\mathcal{O}$ of M, then we have $(\Delta\nu)\big|_{\mathcal{O}} \equiv 0$ and (4.6) implies $(\Delta\phi)\big|_{\mathcal{O}} \equiv c_{\mathcal{O}}$ for a non-zero constant $c_{\mathcal{O}} \in \mathbb{R}$. Thus, (4.8) is satisfied for $F = c_{\mathcal{O}}$ on $\mathcal{O}$. Now we assume that f is non-vanishing on M.

From (1.2), (4.6) and (4.7) we obtain

$$(1,0,0,1) \wedge \zeta = -C, \tag{4.11}$$

where ζ is the vector field given by

$$\zeta = \left(\frac{\Delta\phi_u}{f} e_1 + \frac{\Delta\phi_v}{f} e_2 - \xi e_3 \right). \tag{4.12}$$

From (4.11) we obtain

$$0 = e_i(C) = e_i\big((1,0,0,1) \wedge \zeta\big) = (1,0,0,1) \wedge \widetilde{\nabla}_{e_i}\zeta$$

for $i = 1, 2$ from which we see that $\widetilde{\nabla}_{e_i}\zeta$ is proportional to $(1,0,0,1)$ which is normal to M because of (4.4). Thus, the tangent part of $\widetilde{\nabla}_{e_i}\zeta$ vanishes identically and (4.12) implies

$$\nabla_{e_i}\left(\frac{\Delta\phi_u}{f} e_1 + \frac{\Delta\phi_v}{f} e_2 \right) + \xi A_3(e_i) = 0, \quad i = 1, 2.$$

As $\omega_{12} = 0$, from these equations, (4.3a) and (4.5) we obtain

$$\begin{aligned} \frac{\partial}{\partial u}\left(\frac{\Delta\phi_u}{f} - \phi_u \right) e_1 + \frac{\partial}{\partial u}\left(\frac{\Delta\phi_v}{f} - \phi_v \right) e_2 &= 0 \\ \frac{\partial}{\partial v}\left(\frac{\Delta\phi_u}{f} - \phi_u \right) e_1 + \frac{\partial}{\partial v}\left(\frac{\Delta\phi_v}{f} - \phi_v \right) e_2 &= 0 \end{aligned}$$

from which we have

$$\frac{\Delta\phi_u}{f} - \phi_u = c_1 \quad \text{and} \quad \frac{\Delta\phi_v}{f} - \phi_v = c_2 \tag{4.13}$$

for some constants c_1 and c_2. These equations imply

$$\psi_v \Delta\psi_u - \psi_u \Delta\psi_v = 0, \tag{4.14}$$

where ψ is the function given by (4.8a). By solving (4.14), we obtain (4.8b) for a differentiable function F.

Converse is given in Proposition 4.1. $\square$

5 Marginally trapped surfaces with pointwise 1-type Gauss map

5.1 Marginally trapped surfaces with pointwise 1-type Gauss map of the second kind

Let M be a marginally trapped surface in the Minkowski space-time and $C : M \to \Lambda^{4,2}(\mathbb{E}^4_1) \cong \mathbb{E}^6_3$ be a smooth mapping. We consider an orthonormal frame field $\{e_1, e_2; e_3, e_4\}$ on M. Then, C can be expressed as

$$\begin{aligned} C &= \sum_{1 \leq A < B \leq 4} \varepsilon_A \varepsilon_B C_{AB} e_A \wedge e_B, && (5.1\text{a}) \\ C_{AB} &= \langle C, e_A \wedge e_B \rangle,\ 1 \leq A \leq B \leq 4. && (5.1\text{b}) \end{aligned}$$

Clearly, C is constant if and only if its components satisfy

$$e_i(C_{AB}) = \langle C, \widetilde{\nabla}_{e_i} e_A \wedge e_B + e_A \wedge \widetilde{\nabla}_{e_i} e_B \rangle, \quad 1 \leq A < B \leq 4, \quad i = 1, 2. \tag{5.2}$$

Now, we give the following proposition.

Proposition 5.1. *Let M be an oriented marginally trapped surface in the Minkowski space-time. If M has pointwise 1-type Gauss map of the second kind, then it is pseudo-umbilical.*

Remark 3. In this subsection, the surfaces we are dealing with does not contain any open part with harmonic Gauss map. In expression, for a given surface M, we assume that the interrior of the subset $\{p | (\Delta\nu)\big|_p = 0\}$ of M is empty.

Proof. Suppose that M is a marginally trapped surface in the Minkowski space-time with the Gauss map ν. Then, ν satisfies (3.1) for any local othornormal base field $\{e_1, e_2\}$ of the tangent bundle of M. Now, we assume that M has pointwise 1-type Gauss map of the second kind, i.e., (1.2) is satisfied for a constant non-zero vector C given by (5.1) and a smooth function $f \not\equiv 0$. As C is a constant vector, its components satisfy (5.2). Consider the open subset $\mathcal{F} = \{p \in M | f(p) = 0\}$ and let $\mathcal{O}$ be its interrior. Then we have $\mathcal{O}$ is empty, because of Remark 3.

From (1.2) and (3.1) we obtain

$$-2K\nu + 2K^D e_1 \wedge e_2 + 2\sum_{i=1}^{2} De_i H \wedge e_i = f(\nu + C).$$

From which and (5.1b) we have

$$C_{12} = 2K^D/f, \tag{5.3}$$
$$\langle C, H \wedge e_i \rangle = 0, \quad i = 1, 2 \tag{5.4}$$

on the open subset $\mathcal{M} = M - \mathcal{F}$ of M.

Let $\{e_3, e_4\}$ be an orthonormal basis of the normal bundle of M such that $\varepsilon_3 = -\varepsilon_4 = 1$ and $H = \alpha(e_3 - e_4)$ for a non-vanishing smooth function α. Thus, (5.1b) and (5.4) imply

$$C_{i3} = C_{i4}, \quad i = 1, 2 \tag{5.5}$$

from which we get

$$e_j\left(C_{i3}\right) = e_j\left(C_{i4}\right), \tag{5.6a}$$
$$\langle C, \nabla_{e_j} e_i \wedge (e_3 - e_4) \rangle = 0, \tag{5.6b}$$
$$\langle C, e_i \wedge D_{e_j}(e_3 - e_4) \rangle = 0, \quad i, j = 1, 2 \tag{5.6c}$$

on the open set $\mathcal{M}$. By a simple calculation using (5.2) and (5.6a), we get

$$\langle C, \widetilde{\nabla}_{e_j} e_i \wedge e_3 + e_i \wedge \widetilde{\nabla}_{e_j} e_3 \rangle = \langle C, \widetilde{\nabla}_{e_j} e_i \wedge e_4 + e_i \wedge \widetilde{\nabla}_{e_j} e_4 \rangle. \tag{5.7}$$

By taking into account (2.1) and (2.2), we see that (5.7) implies

$$\begin{gathered}\langle C, \nabla_{e_j} e_i \wedge e_3 + h(e_i, e_j) \wedge e_3 - e_i \wedge A_3 e_j + e_i \wedge D_{e_j} e_3 \rangle = \\ \langle C, \nabla_{e_j} e_i \wedge e_4 + h(e_i, e_j) \wedge e_4 - e_i \wedge A_4 e_j + e_i \wedge D_{e_j} e_4 \rangle.\end{gathered}$$

From this equation, (5.1b), (5.6b) and (5.6c) we obtain

$$h^4_{ij} C_{34} - h^3_{j(3-i)} C_{i(3-i)} = h^3_{ij} C_{34} - h^4_{j(3-i)} C_{i(3-i)}, \ i, j = 1, 2$$

on $\mathcal{M}$. Therefore, the functions $C_{12} = -C_{21},\ C_{34},\ h^3_{11},\ h^3_{12}, \ldots, h^4_{22}$ satisfy

$$\mathbf{A} \begin{pmatrix} C_{12} \\ C_{34} \end{pmatrix} = 0 \tag{5.8}$$

for a 4×2 matrix $\mathbf{A}$ given by

$$\mathbf{A} = \begin{pmatrix} h^3_{11} - h^4_{11} & -h^3_{12} + h^4_{12} \\ h^3_{12} - h^4_{12} & h^3_{11} - h^4_{11} \\ h^3_{12} - h^4_{12} & -h^3_{22} + h^4_{22} \\ h^3_{22} - h^4_{22} & h^3_{12} - h^4_{12} \end{pmatrix}.$$

Note that rank$\mathbf{A} < 2$ if and only if $A_3 = A_4$.

Now we will show that $A_3 = A_4$. Suppose $A_3 \neq A_4$ at a point p of $\mathcal{M}$. Then, from (5.8) we see that there exists a neighborhood $\mathcal{N}_p$ of p on $\mathcal{M}$ such that $C_{34} = C_{12} \equiv 0$ on $\mathcal{N}_p$. Thus, (5.3) implies $K^D \equiv 0$ on $\mathcal{N}_p$, i.e. the open part $\mathcal{N}_p$ of M has flat normal bundle. Thus, there exist an orthonormal base field $\{e_1, e_2\}$ of tangent bundle of $\mathcal{N}_p$ such that

$$A_\beta = \mathrm{diag}(h_1^\beta, h_2^\beta), \quad \beta = 3, 4 \tag{5.9}$$

for some smooth functions h_1^β, h_2^β satisfying

$$h_1^3 + h_2^3 = h_1^4 + h_2^4 = 2\alpha. \tag{5.10}$$

On the other hand, we have $e_i(C_{34}) = 0,\ i = 1, 2$ on $\mathcal{N}_p$, as $C_{34}\big|_{\mathcal{N}_p} = 0$. By using these equations, (2.2) and (5.2), we obtain $e_i(C_{34}) = \langle C, -(A_3 e_i) \wedge e_4 + (A_4 e_i) \wedge e_3 \rangle = 0,\ i = 1, 2$ from which, (5.5) and (5.9) we get

$$(h_1^3 - h_1^4) C_{13} = (h_2^3 - h_2^4) C_{23} = 0$$

on $\mathcal{N}_p$. As $C \neq 0$, these equations and (5.10) imply $h_1^3 - h_1^4 = h_2^3 - h_2^4 \equiv 0$. Thus, $A_3 = A_4$ over $\mathcal{N}_p$ which yields a contradiction. Therefore, we have proved $A_3 = A_4$ on $\mathcal{M}$.

Since $A_3 = A_4$ on $\mathcal{M}$, we have $A_H\Big|_{\mathcal{M}} = 0$. Because $\mathcal{O}$ is empty, and the functions $\zeta_{ij} = \langle A_H e_i, e_j \rangle$ are continous, we have $A_H = 0$ over M. $\square$

By combaining Proposition 4.2, Proposition 5.1 and Remark 2, we state the following classification theorem.

Theorem 5.2. *Let M be a marginally trapped surface in the Minkowski space-time and $\mathcal{M}$. Then, M has pointwise 1-type Gauss map of the second kind if and only if it is congruent to the surface given in Proposition 4.1.*

By combaining Proposition 3.3 and Theorem 5.2, we have the following classification theorem:

Theorem 5.3. *Let M be a marginally trapped surface in the Minkowski space-time and $\mathcal{M}$ the open subset of M given by $\mathcal{M} = \{p \in M | \Delta\phi|_p \neq 0\}$. Then, M has pointwise 1-type Gauss map if and only if $\mathcal{M}$ is union of these type of surfaces.*

(i) a non-flat, non-parallel surface lying in the de Sitter space-time $S_1^3(r^2)$ for some $r > 0$ such that the mean curvature vector H' of M in $S_1^3(r^2)$ satisfies $\langle H', H' \rangle = -r^2$;

(ii) a non-flat, non-parallel surface lying in the hyperbolic space $H^3(-r^2)$ *for some* $r > 0$ *such that the mean curvature vector* H' *of* M *in* $H^3(-r^2)$ *satisfies* $\langle H', H'\rangle = r^2$*;*

(iii) a surface congruent to the surface given in Proposition 4.1.

By combaining Proposition 4.1 and Theorem 5.3, we obtain classification of marginally trapped surfaces with 1-type Gauss map.

Proposition 5.4. *Let* M *be a marginally trapped surface in the Minkowski space-time. Then,* M *has non-harmonic 1-type Gauss map, if and only if it is locally congruent to the surface given by* (4.2) *for a smooth function* $\phi : \Omega \to \mathbb{R}$ *satisfying Helmholtz equation*

$$\Delta\phi + \lambda\phi = c_1 u + c_2 v \tag{5.11}$$

for some constants $\lambda \neq 0,\ c_1,\ c_2$.

A surface is said to be null 2-type if its position vector x can be expressed as a sum $x = x_0 + x_1$ for some non-constant mappings x_0 and x_1 such that $\Delta x_0 = 0$ and $\Delta x_1 = \lambda x_1,\ \lambda \in \mathbb{R} - \{0\}$. Next, we want to give the following corollary of Proposition 5.4.

Corollary 5.5. *If a marginally trapped surface* M *has non-harmonic, 1-type Gauss map, then it is null 2-type.*

Proof. We assume that M is the surface given by (4.2) for a smooth function ϕ satisfying (5.11). We define the mappings x_0 and x_1 as

$$x_0(u,v) = \Big(-\frac{\psi}{\lambda}, u, v, -\frac{\psi}{\lambda}\Big)$$

and

$$x_1(u,v) = \Big(\phi + \frac{\psi}{\lambda}, 0, 0, \phi + \frac{\psi}{\lambda}\Big).$$

We see that these mappings satisfy $x = x_0 + x_1$, $\Delta x_0 = 0$ and $\Delta x_1 = \lambda x_1$. Hence M is null 2-type. □

5.2 Examples

In this subsection we present explicit examples of (partly) marginally trapped surfaces with pointwise 1-type Gauss map. We also give construction of marginally trapped surfaces with 1-type Gauss map for a given boundary curve.

In the first example, we construct a marginally trapped surface with proper pointwise 1-type Gauss map of the second kind.

Example 1. Let $\epsilon > 0$ and $\Omega_\epsilon = \{(u, v) | \epsilon < u + v < 1\}$ and $\phi : \Omega \to \mathbb{R}$ be a function given by

$$\phi(u, v) = \exp\left(\frac{1}{u + v}\right)$$

Then, by a direct calculation, we see that $\phi : \Omega_\epsilon \to \mathbb{R}$ satisfies (4.8) for the differentiable function

$$F(\phi) = \big(- 2(\ln \phi)^4 - 4(\ln \phi)^3\big)\phi \tag{5.12}$$

is differentiable and the function given by (4.9) becomes

$$f(u, v) = -\frac{12}{(u + v)^2} - \frac{12}{(u + v)^3} - \frac{2}{(u + v)^4} \tag{5.13}$$

which is smooth on Ω_ϵ. Hence, Proposition 4.1 implies that the marginally trapped surface M given by (4.2) has pointwise 1-type Gauss map of the second kind.

In the next example, we construct a marginally trapped surface with the boundary of a square lying in the space-like plane $\Pi = \{(0, u, v, 0) | u, v \in \mathbb{R}\}$.

Example 2. The function $\phi(u, v) = \sin(n\pi x)sin(n\pi y)$ solves the boundary value problem (2.7) for $\Omega = (0, 1) \times (0, 1)$, where $n \in \mathbb{N}$. Therefore, the surface given by $x = (\phi(u, v), u, v, \phi(u, v))$ has 1-type Gauss map. Moreover, the boundary curve of this surface is the non-smooth curve β which is a square in the plane Π.

In the next proposition, we give a generalization of Example 2 and obtain construction of marginally trapped surfaces in the Minkowski space-time with 1-type Gauss map for a given (smooth or non-smooth) boundary curve.

Proposition 5.6. *Let β be a continuous Jordan curve lying in a space-like plane of the Minkowski space-time. Then, for each eigenvalue λ_k of operator Δ given in* (2.6) *there exists a (partly) marginally trapped surface M_k with the boundary β and 1-type Gauss map satisfying* (1.2) *for $f \equiv \lambda_k$ and* $C = (1, 0, 0, 1) \wedge (0, 0, 0, 1)$.

Proof. Up to isometries of the Minkowski space-time, we assume that β is contained by the space-like plane $\Pi = \{(0, u, v, 0) | u, v \in \mathbb{R}\}$. Let Ω be the open subset of Π bounded by β. As β is a Jordan curve, Ω is connected. Consider a solution ϕ_k of the boundary value problem given in (2.7) for an eigenvalue λ_k of (2.6). Then, ϕ_k is smooth (see Section 2.3) and satisfies (5.11) for $c_1 = c_2 = 0$.

Let M_k be (partly) marginally trapped surface in the Minkowski space-time given by (4.2) for $\phi = \phi_k$. Then, ϕ satisfies (4.8) for the function $F = \lambda_k \phi$. Moreover, the function f given by (4.8) becomes a constant function. Thus, it is smooth. Hence, M_k has 1-type Gauss map because of Proposition 4.1. Moreover, as $\phi_k = 0$ on $\partial\Omega = \beta$, β is the boundary curve of M_k. $\square$

In the next example, we construct a partly marginally trapped surface with 1-type Gauss map whose boundary curve is smooth.

Example 3. Consider the function defined by $\phi : \Omega \to \mathbb{R}$

$$\phi(u,v) = \begin{cases} \frac{J_1\left(k\sqrt{u^2+v^2}\right)u}{\sqrt{u^2+v^2}} & \text{if} \quad 0 < u^2+v^2 < 1 \\ 0 & \text{if} \quad u^2+v^2 = 0, \end{cases}$$

where J_1 denotes the Bessel function of the first kind of order 1, given by

$$J_1(t) = \sum_{i=0}^{\infty} \frac{(-\frac{t}{2})^{2i+1}}{i!(i+1)!},$$

$\Omega = \{(u,v) \mid u^2+v^2 < 1\}$ and k is the first zero of J_1. Then, ϕ satisfies (5.11) for $\lambda = 1,\ c_1 = c_2 = 0$. Thus, the surface given by $x = (\phi(u,v), u, v, \phi(u,v))$ has 1-type Gauss map. Moreover, the boundary curve of this surface is a circle in the plane Π. M is partly marginally trapped because $H = 0$ at $p = x(0,0)$. Note that the function ϕ is obtained by using a solution of (2.7) for the given Ω.

References

[1] D. D. Bleecker, J. L. Weiner *Extrinsic bounds on λ_1 of Δ on a compact manifold.* Comment Math. Helv. vol. 51(1976) 601-609.

[2] J. L. Cabrerizo, M. Fernández and J. S. Gómez, *Isotropy and marginally trapped surfaces in a spacetime.* Class. Quantum Grav. 27 (2010) 135005 (12pp).

[3] B. Y. Chen, *Total mean curvature and submanifolds of finite type*, World Scientific, Singapor-New Jersey-London, 1984.

[4] B. Y. Chen, *Finite type pseudo-Riemannian submanifolds.* Tamkang J. of Math, vol. 17, no 2 (1986), 137-151.

[5] B. Y. Chen, *A report on submanifolds of finite type.* Soochow J. Math., vol. 22 (1996), 117-337.

[6] B. Y. Chen, *Pseudo-Riemannian geometry, δ-invariants and applications*, World Scientific, Singapor-New Jersey-London, 2011.

[7] B.Y. Chen and S. Ishikawa, *Biharmonic surfaces in pseudo-Euclidean spaces*, Mem. Fac. Sci. Kyushu Univ. Ser. A 45 (1991), no. 2, 323-347.

[8] B. Y. Chen, J. M. Morvan and T. Nore , *Energy, tension and finite type maps.* Kodai Math. J. vol. 9, num 3(1986) 406-418.

[9] B.Y. Chen and P. Piccinni, *Submanifolds with finite type Gauss map*, Bull. Austral. Math. Soc., **35** (1987), 161–186.

[10] B.Y. Chen and J. van der Veken, *Marginally trapped surfaces in Lorentzian space forms with positive relative nullity.* Class. Quantum Grav. 24 (2007) 551–563

[11] B. Y. Chen and J. van der Veken, *Classification of marginally trapped surfaces with parallel mean curvature vector in Lorentzian space forms.* Houston J. Math. 36 (2010), no. 2, 421-449.

[12] U. Dursun, *Hypersurfaces with pointwise 1-type Gauss map*, Taiwanese J. Math., 11(2007), 1407–1416.

[13] U. Dursun, *Hypersurfaces with pointwise 1-type Gauss map in Lorentz-Minkowski space*, Proc. Est. Acad. Sci., 58(2009), 146–161.

[14] U. Dursun and N. C. Turgay, *Space-like Surfaces in Minkowski Space $\mathbb{E}^4_1$ with Pointwise 1-Type Gauss Map.* (submitted).

[15] L. R. Evans, *Partial differential equations.* American Mathematical Society, Providence, RI, 1998.

[16] S. Haesen and M. Ortega *Boost invariant marginally trapped surfaces in Minkowski 4-space* Class. Quantum Grav. 24 (2007) 5441.

[17] Y. H. Kim and Y. W. Yoon, *Classification of Rotation Surfaces in Pseudo-Euclidean Space*, J. Korean Math., 41(2004), 379–396.

[18] R. Penrose, *Gravitational collapse and space-time singularities.* Phys. Rev. Lett. 14 (1965), 57–59.

[19] M. P. O'Neill, *Semi-Riemannian geometry with applications to relativity.* Academic Press, 1983.

[20] R. C. Reilly, *On the first eigenvalue of the Laplacian for compact submanifolds of Euclidean space.*. Comment. Math. Helv. 52(1977), Issue 1, 525–533.

[21] J. Senovilla, *Trapped surfaces, horizons and exact solutions in higher dimensions.* Class. Quantum Grav. 19 (2002), L113–L119.

[22] Y. W. Yoon, *On the Gauss map of translation surfaces in Minkowski 3-spaces*, Taiwanese J. Math., 6(2002), 389–398.